

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 June 2002 (20.06.2002)

PCT

(10) International Publication Number
WO 02/48297 A1

(51) International Patent Classification⁷: **C11D 1/86 //**
1:14, 1:22, 1:62, 1:72

Sunlight, Quarry Road East, Bebington, Wirral, Mersey-
side CH63 3JW (GB).

(21) International Application Number: **PCT/EP01/14488**

(74) Agents: **FRANSELLA, Mary, Evelyn et al.**; Unilever
PLC, Patent Dept., Colworth House, Sharnbrook, Bedford,
Bedfordshire MK44 1LQ (GB).

(22) International Filing Date: 6 December 2001 (06.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0030669.6 15 December 2000 (15.12.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZW.

(71) Applicant (*for AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH,
GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SL, SZ,
TT, TZ, UG, ZA, ZW only*): **UNILEVER PLC** [GB/GB];
Unilever House, Blackfriars, London EC4P 4BQ (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(71) Applicant (*for all designated States except AE, AG, AU,
BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, IN, KE, LC,
LK, LS, MN, MW, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA,
ZW*): **UNILEVER N.V.** [NL/NL]; Weena 455, NL-3013
AL Rotterdam (NL).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- entirely in electronic form (except for this front page) and
available upon request from the International Bureau

(71) Applicant (*for IN only*): **HINDUSTAN LEVER LIM-
ITED** [IN/IN]; Hindustan Lever House, 165/166 Backbay
Reclamation, Maharashtra, 400 020 Mumbai (IN).

(72) Inventors: **SINGH, Amrat, Paul**; 8 Kingsbrook Way,
Higher Bebington, Wirral, Merseyside CH63 5NG (GB).
THORLEY, David, Christopher; Unilever Research Port

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/48297 A1

(54) Title: **DETERGENT COMPOSITIONS**

(57) Abstract: A built laundry detergent composition contains a combination of anionic, nonionic and cationic surfactants which includes a highly ethoxylated nonionic surfactant which is a C8-C16 alcohol ethoxylated with an average of from 20 to 50 ethylene oxide groups. The cationic surfactant may advantageously be an ethoxylated quaternary ammonium compound. The composition exhibits improved oily soil detergency.

- 1 -

DETERGENT COMPOSITIONSTECHNICAL FIELD

5

The present invention relates to laundry detergent compositions containing a combination of anionic, specified nonionic and cationic surfactants giving improved oily soil detergency.

10

BACKGROUND

Heavy duty laundry detergent compositions have for many years contained anionic sulphonate or sulphate surfactant, for example, linear alkylbenzene sulphonate (LAS), together with ethoxylated nonionic surfactants. Examples abound in the published literature.

The preferred ethoxylated alcohol nonionic surfactants giving an optimum balance of properties have generally been those having an alkyl chain length of C₁₂-C₁₅ and an average degree of ethoxylation of 1 to 10, preferably 3 to 7, more preferably about 5.

Longer-chain alcohols having a higher degree of ethoxylation, for example, tallow alcohol (C₁₈) 11EO, have also been used.

These relatively hydrophobic materials of low HLB value are generally liquids at ambient temperature and exhibit excellent oily soil removal.

- 2 -

Longer-chain alcohols having higher degrees of ethoxylation, for example, tallow (C₁₈) alcohol 25EO and 50EO, are solids at ambient temperature and are used as slowly dissolving coating materials, for example, for enzyme or antifoam
5 granules.

Laundry detergent compositions containing cationic (quaternary ammonium) surfactants in combination with anionic and nonionic surfactants are widely disclosed in the
10 patent literature.

It has now surprisingly been found that the combination of anionic and cationic surfactants together with ethoxylated alcohols combining a shorter alkyl chain length and a higher
15 degree of ethoxylation can give enhanced oily soil removal.

PRIOR ART

- 20 WO 94 16052A (Unilever) discloses high bulk density laundry powders based on LAS and conventional nonionic surfactants, and containing small amounts of very highly ethoxylated alcohols, eg tallow alcohol 80EO, as a dissolution aid.
- 25 WO 93 02176A (Henkel) discloses the use of highly ethoxylated aliphatic alcohols as "structure breakers" in high bulk density powders containing conventional nonionic surfactants.
- 30 EP 293 139A (Procter & Gamble) discloses twin-compartment sachets containing detergent powders. Some powders contain very small amounts of tallow alcohol 25EO.

- 3 -

US 4 294 711 (Procter & Gamble) discloses a textile softening heavy duty detergent composition containing 1 wt% of tallow alcohol 80EO.

- 5 WO 92 18594A (Procter & Gamble) discloses builder granules of layered silicate coated with tallow alcohol 50EO.

EP 142 910A and EP 495 345A (Procter & Gamble) disclose antifoam granules containing highly ethoxylated
10 alcohols.

WO 93 19148A (Procter & Gamble) discloses liquid hard surface cleaning compositions containing highly ethoxylated nonionic surfactants optionally plus anionic surfactant.

15

WO 97 43364, WO 97 43365A, WO 97 43371A, WO 97 43387A, WO 97 43389A, WO 97 43390A, WO 97 43391A and WO 97 43393A, (Procter & Gamble) disclose laundry detergent compositions containing so-called "AQA" alkoxylated quaternary ammonium
20 surfactants in combination with other surfactants.

DEFINITION OF THE INVENTION

25 The present invention provides a built laundry detergent composition comprising


(i) from 5 to 40 wt%, preferably from 7 to 30 wt%, of surfactant consisting essentially of:

30

(i) (a) an anionic sulphonate or sulphate surfactant,

- 4 -

(i) (b) an ethoxylated alcohol nonionic surfactant of the general formula I



5

wherein R is a hydrocarbyl chain having from 8 to 16 carbon atoms, and the average degree of ethoxylation n is from 20 to 50,

10 (i) (c) a quaternary ammonium cationic surfactant,

(ii) from 10 to 80 wt% of detergency builder,

(iii) optionally other detergent ingredients to 100 wt%.

15


The invention also provides a process for laundering textile fabrics by machine or hand, which includes the step of immersing the fabrics in a wash liquor comprising water in which a laundry detergent composition as defined in the
20 previous paragraph is dissolved or dispersed.

The invention further provides the use of a surfactant (i) consisting essentially of

25 (i) (a) an anionic sulphonate or sulphate surfactant,

(i) (b) an ethoxylated alcohol nonionic surfactant of the general formula I

30



- 5 -

wherein R is a hydrocarbyl chain having from 8 to 16 carbon atoms, and the average degree of ethoxylation n is from 20 to 50,

- 5 (i) (c) a quaternary ammonium cationic surfactant,
in a laundry detergent composition in an amount of from 5 to 40 wt%, to improve the oily soil detergency of the composition.

10

DETAILED DESCRIPTION OF THE INVENTION

The detergent compositions of the invention contain a
15 combination of an anionic sulphonate or sulphate surfactant, a defined nonionic surfactant, and a cationic surfactant. The total amount of the three surfactants is from 5 to 40 wt%, preferably from 7 to 30 wt%.

20 Detergent compositions according to the invention show improved oily soil detergency across a range of fabrics and water hardnesses.

25 The anionic surfactant (i) (a)

The anionic surfactant is a sulphonate or sulphate anionic surfactant.

30 Anionic surfactants are well-known to those skilled in the art. Many suitable detergent-active compounds are available and are fully described in the literature, for example, in

- 6 -

"Surface-Active Agents and Detergents", Volumes I and II, by Schwartz, Perry and Berch.

Examples include alkylbenzene sulphonates, primary and
5 secondary alkylsulphates, particularly C₈-C₁₅ primary alkyl
sulphates; alkyl ether sulphates; olefin sulphonates;
alkyl xylene sulphonates; dialkyl
sulphosuccinates; and fatty acid ester sulphonates.
Sodium salts are generally preferred.

10


Preferably the anionic surfactant is linear alkylbenzene
sulphonate or primary alcohol sulphate. More preferably the
anionic surfactant is linear alkylbenzene sulphonate.

15

The ethoxylated nonionic surfactant (i) (b)

The nonionic surfactant is an ethoxylated aliphatic alcohol
of the formula

20



wherein R is a hydrocarbyl chain having from 8 to 16 carbon
atoms, and the average degree of ethoxylation n is from 20
25 to 50.

The hydrocarbyl chain, which is preferably saturated,
preferably contains from 10 to 16 carbon atoms, more
preferably from 12 to 15 carbon atoms. In commercial
30 materials containing a spread of chain lengths, these
figures represent an average.

- 7 -

The alcohol may be derived from natural or synthetic feedstock. Preferred alcohol feedstocks are coconut, predominantly C₁₂-C₁₄, and oxo C₁₂-C₁₅ alcohols. Longer chain materials such as tallow or hardened tallow
5 (C₁₈) are not preferred.

The average degree of ethoxylation ranges from 20 to 50, preferably from 25 to 40.

10 Preferred materials have an average alkyl chain length of C₁₂-C₁₆ and an average degree of ethoxylation of 25 to 40.

An example of a suitable commercially available material is Lutensol (Trade Mark) AO30, ex BASF, which is a C₁₃-C₁₅
15 alcohol having an average degree of ethoxylation of 30.

The cationic surfactant (i) (c)


20 Preferred water-soluble cationic surfactants are quaternary ammonium salts of the general formula II



25 wherein R₁ is a relatively long (C₈-C₁₈) hydrocarbon chain, typically an alkyl, hydroxyalkyl or ethoxylated alkyl group, optionally interrupted with a heteroatom or an ester or amide group; each of R₂, R₃ and R₄ (which may be the same or different) is a short-chain (C₁-C₃) alkyl or substituted
30 alkyl group; and X is a solubilising anion, for example a chloride, bromide or methosulphate ion.


- 8 -

According to a first preferred embodiment of the invention, the cationic surfactant is a quaternary ammonium compound of the formula II in which R_1 is a C_8 - C_{18} alkyl group, more preferably a C_8 - C_{10} or C_{12} - C_{14} alkyl group, R_2 is a methyl group, and R_3 and R_4 , which may be the same or different, are methyl or hydroxyethyl groups. Such compounds have the formula III:



In an especially preferred compound, R_1 is a C_{12} - C_{14} alkyl group, R_2 and R_3 are methyl groups, R_4 is a 2-hydroxyethyl group, and X^- is a chloride ion. This material is available commercially as Praepagen (Trade Mark) HY from Clariant GmbH, in the form of a 40 wt% aqueous solution.

According to a second preferred embodiment of the invention, the cationic surfactant is an ethoxylated quaternary ammonium compound of the formula IV:



wherein R_5 is a C_6 - C_{20} alkyl group,
 m is an integer from 1 to 20,

- 9 -

R_6 and R_7 , which may be the same or different, each represents a C_1 - C_4 alkyl group or a C_2 - C_4 hydroxyalkyl group,

R_8 represents a C_1 - C_4 alkyl group, and

5 Y^- represents a monovalent solubilising anion.


In preferred ethoxylated cationic surfactants of the formula IV used in accordance with the invention,

10 R_5 is a C_{10} - C_{16} alkyl group,
 m is from 1 to 4,
 R_6 , R_7 and R_8 are methyl groups, and
 Y^- represents Cl^- .

15 An especially preferred ethoxylated cationic surfactant used in accordance with the present invention is of the formula IV in which

R_5 is a C_{12} - C_{14} alkyl group,
 m is 3,
 R_6 , R_7 and R_8 are methyl groups, and
 20 Y^- represents Cl^- .

This material (DBETAC) has the formula V



30 wherein R_9 is a C_{12} - C_{14} alkyl group.

Other classes of cationic surfactant include cationic esters (for example, choline esters).

- 10 -

The surfactant combination (i)

The surfactant combination preferably consists essentially of:

5

(i) (a) from 20 to 98 wt% of the anionic sulphonate or sulphate detergent;

10

(i) (b) from 1 to 60 wt% of the nonionic surfactant, and

(i) (c) from 1 to 60 wt% of the cationic surfactant.

The whole product preferably contains from:

15 (i) (a) from 1 to 20 wt% of the anionic sulphonate or sulphate detergent;

(i) (b) from 0.5 to 20 wt% of the nonionic surfactant, and

20 (i) (c) from 0.1 to 20 wt% of the cationic surfactant.

Preferred weight ratios are as follows:

(i) (a) : (i) (b)	1.5:1 - 25:1
(i) (a) : (i) (c)	0.2:1 - 5:1
(i) (b) : (i) (c)	0.1:1 - 3:1

25

- 11 -

In the first preferred embodiment of the invention wherein the cationic surfactant is a compound of the formula III, then the surfactant system is preferably composed as follows:

5

(i) (a) from 50 to 98 wt%, preferably from 60 to 95 wt%, of the anionic sulphonate or sulphate detergent;

(i) (b) from 1 to 30 wt%, preferably from 5 to 25 wt%, of the nonionic surfactant, and

10

(i) (c) from 1 to 30 wt% , preferably from 5 to 25 wt%, of the cationic surfactant.

The whole product preferably contains:

15

(i) (a) from 3 to 30 wt%, preferably from 5 to 25 wt%, of the anionic sulphonate or sulphate detergent;

(i) (b) from 0.5 to 10 wt%, preferably from 1 to 5 wt%, of the nonionic surfactant, and

20

(i) (c) from 0.1 to 10 wt%, preferably from 0.2 to 5 wt%, of the cationic surfactant.

25 Preferred weight ratios are as follows:

	Preferred	Most preferred
(i) (a) : (i) (b)	2:1 - 25:1	3:1 - 20:1
(i) (a) : (i) (c)	2:1 - 50:1	3:1 - 30:1
(i) (b) : (i) (c)	0.1:1 - 3:1	0.5:1 - 2:1

- 12 -

In the second preferred embodiment of the invention wherein the cationic surfactant is an ethoxylated compound of the formula IV, then the surfactant system is preferably composed as follows:

5

(i) (a) from 20 to 80 wt%, preferably from 30 to 60 wt%, of the anionic sulphonate or sulphate detergent;

(i) (b) from 5 to 40 wt%, preferably from 10 to 30 wt%, of
10 the nonionic surfactant, and

(i) (c) from 10 to 60 wt% , preferably from 20 to 50 wt%, of the cationic surfactant.

15 The whole product preferably contains:

(i) (a) from 3 to 30 wt%, preferably from 5 to 25 wt%, of the anionic sulphonate or sulphate detergent;

20 (i) (b) from 0.5 to 10 wt%, preferably from 1 to 5 wt%, of the nonionic surfactant, and

(i) (c) from 0.1 to 10 wt%, preferably from 0.2 to 5 wt%, of the cationic surfactant.

25

- 13 -

Preferred weight ratios are as follows:

	Preferred	Most preferred
(i) (a) : (i) (b)	1.5:1 - 10:1	1:1 - 5:1
(i) (a) : (i) (c)	0.2:1 - 5:1	0.5:1 - 3:1
(i) (b) : (i) (c)	0.2:1 - 5:1	0.5:1 - 3:1

- 5 Optionally minor, non-interfering amounts of other surfactants may also be present. Preferably, however, the composition is free from nonionic surfactants other than the defined nonionic surfactant (i) (b).
- 10 More preferably the composition is substantially free of other non-soap surfactants.

Optionally soap may also be present, for example, in an amount of from 1 to 5 wt%.

15

Detergency builder (ii)

- The compositions may suitably contain from 10 to 80%, preferably from 15 to 70% by weight, of detergency builder. Preferably, the quantity of builder is in the range of from 15 to 50% by weight.
- 20

- Preferably the builder is selected from sodium tripolyphosphate, zeolite, sodium carbonate, sodium citrate, layered silicate, and combinations of these.
- 25

- 14 -

The zeolite used as a builder may be the commercially available zeolite A (zeolite 4A) now widely used in laundry detergent powders. Alternatively, the zeolite may be maximum aluminium zeolite P (zeolite MAP) as described and claimed
5 in EP 384 070B (Unilever), and commercially available as Doucil (Trade Mark) A24 from Ineos Silicas Ltd, UK. Zeolite MAP is defined as an alkali metal aluminosilicate of zeolite P type having a silicon to aluminium ratio not exceeding 1.33, preferably within the range of from 0.90 to
10 1.33, preferably within the range of from 0.90 to 1.20.

Especially preferred is zeolite MAP having a silicon to aluminium ratio not exceeding 1.07, more preferably about 1.00. The particle size of the zeolite is not critical.
15 Zeolite A or zeolite MAP of any suitable particle size may be used.

Also preferred according to the present invention are phosphate builders, especially sodium tripolyphosphate.
20 This may be used in combination with sodium orthophosphate, and/or sodium pyrophosphate.

Other inorganic builders that may be present additionally or alternatively include sodium carbonate, layered silicate,
25 amorphous aluminosilicates.

Organic builders that may be present include polycarboxylate polymers such as polyacrylates and acrylic/maleic copolymers; polyaspartates; monomeric polycarboxylates such
30 as citrates, gluconates, oxydisuccinates, glycerol mono-di- and trisuccinates, carboxymethyloxysuccinates, carboxy-methyloxymalonates, dipicolinates,

- 15 -

hydroxyethyliminodiacetates, alkyl- and alkenylmalonates and succinates; and sulphonated fatty acid salts.

Organic builders may be used in minor amounts as supplements
5 to inorganic builders such as phosphates and zeolites.
Especially preferred supplementary organic builders are citrates, suitably used in amounts of from 5 to 30 wt%, preferably from 10 to 25 wt%; and acrylic polymers, more especially acrylic/maleic copolymers, suitably used in
10 amounts of from 0.5 to 15 wt%, preferably from 1 to 10 wt%.

Builders, both inorganic and organic, are preferably present in alkali metal salt, especially sodium salt, form.

15

Other detergent ingredients

As well as the surfactants and builders discussed above, the compositions may optionally contain bleaching components and
20 other active ingredients to enhance performance and properties.

These optional ingredients may include, but are not limited to, any one or more of the following: soap, peroxyacid and
25 persalt bleaches, bleach activators, sequestrants, cellulose ethers and esters, other antiredeposition agents, sodium sulphate, sodium silicate, sodium chloride, calcium chloride, sodium bicarbonate, other inorganic salts, fluorescers, photobleaches, polyvinyl pyrrolidone, other dye
30 transfer inhibiting polymers, foam controllers, foam boosters, acrylic and acrylic/maleic polymers, proteases, lipases, cellulases, amylases, other detergent enzymes,

- 16 -

citric acid, soil release polymers, fabric conditioning compounds, coloured speckles, and perfume.

Detergent compositions according to the invention may
5 suitably contain a bleach system. The bleach system is preferably based on peroxy bleach compounds, for example, inorganic persalts or organic peroxyacids, capable of yielding hydrogen peroxide in aqueous solution. Suitable peroxy bleach compounds include organic peroxides such as
10 urea peroxide, and inorganic persalts such as the alkali metal perborates, percarbonates, perphosphates, persilicates and persulphates. Preferred inorganic persalts are sodium perborate monohydrate and tetrahydrate, and sodium percarbonate. Especially preferred is sodium percarbonate
15 having a protective coating against destabilisation by moisture. Sodium percarbonate having a protective coating comprising sodium metaborate and sodium silicate is disclosed in GB 2 123 044B (Kao).

20 The peroxy bleach compound is suitably present in an amount of from 5 to 35 wt%, preferably from 10 to 25 wt%.

The peroxy bleach compound may be used in conjunction with a bleach activator (bleach precursor) to improve bleaching
25 action at low wash temperatures. The bleach precursor is suitably present in an amount of from 1 to 8 wt%, preferably from 2 to 5 wt%.

Preferred bleach precursors are peroxycarboxylic acid
30 precursors, more especially peracetic acid precursors and peroxybenzoic acid precursors; and peroxycarbonic acid precursors. An especially preferred bleach precursor

- 17 -

suitable for use in the present invention is N,N,N',N'-tetracetyl ethylenediamine (TAED). Also of interest are peroxybenzoic acid precursors, in particular, N,N,N-trimethylammonium toluoyloxy benzene sulphonate.

5

A bleach stabiliser (heavy metal sequestrant) may also be present. Suitable bleach stabilisers include ethylenediamine tetraacetate (EDTA) and the polyphosphonates such as Dequest (Trade Mark), EDTMP.

10

The detergent compositions may also contain one or more enzymes. Suitable enzymes include the proteases, amylases, cellulases, oxidases, peroxidases and lipases usable for incorporation in detergent compositions.

15

In particulate detergent compositions, detergency enzymes are commonly employed in granular form in amounts of from about 0.1 to about 3.0 wt%. However, any suitable physical form of enzyme may be used in any effective amount.

20

Antiredeposition agents, for example cellulose esters and ethers, for example sodium carboxymethyl cellulose, may also be present.

25

The compositions may also contain soil release polymers, for example sulphonated and unsulphonated PET/POET polymers, both end-capped and non-end-capped, and polyethylene glycol/polyvinyl alcohol graft copolymers such as Sokolan (Trade Mark) HP22. Especially preferred soil release

30

polymers are the sulphonated non-end-capped polyesters described and claimed in WO 95 32997A (Rhodia Chimie).

- 18 -

Product form and preparation

The compositions of the invention may be of any suitable physical form, for example, particulates (powders, granules, tablets), liquids, pastes, gels or bars.

According to one especially preferred embodiment of the invention, the detergent composition is in particulate form.

10 Powders of low to moderate bulk density may be prepared by spray-drying a slurry, and optionally postdosing (dry-mixing) further ingredients. "Concentrated" or "compact" powders may be prepared by mixing and granulating processes, for example, using a high-speed mixer/granulator, or other
15 non-tower processes.

Tablets may be prepared by compacting powders, especially "concentrated" powders.

20 Also preferred are liquid detergent compositions, which may be prepared by admixing the essential and optional ingredients in any desired order to provide compositions containing the ingredients in the the requisite concentrations.

25

EXAMPLES

The invention is illustrated in further detail by the
30 following non-limiting Examples, in which parts and percentages are by weight unless otherwise stated.

- 19 -

EXAMPLES 1 to 8, COMPARATIVE EXAMPLES A to NPerformance appraisal of anionic/nonionic/cationic
surfactant mixtures on various soils

- 5 Surfactant mixtures were prepared by mixing sodium linear alkylbenzene sulphonate (LAS), the ethoxylated nonionic surfactant Lutensol AO30 (R = C₁₂-C₁₅ alkyl, n has an average value of 30), and the cationic surfactant Praepagen HY (C₁₂-C₁₄ alkyl methyl hydroxyethyl ammonium chloride) in
- 10 various proportions. High suds detergent compositions suitable for machine or handwash were prepared to the following general formulation:

Total surfactant (LAS + Lutensol AO30 + Praepagen HY)	24.00
Sodium tripolyphosphate	14.50
Sodium carboxymethyl cellulose	0.33
Sodium neutral silicate	6.98
Sodium sulphate	17.75
Fluorescers	0.19
Acrylic/maleic copolymer	1.50
Sodium carbonate	15.00
Sodium perborate monohydrate	8.00
Tetracetyl ethylenediamine	2.40
Phosphonate sequestrant	0.40
Enzyme granules	0.91
Antifoam granules	-
Soil release polymer	0.80
Perfume	0.30
Miscellaneous salts, water etc	to 100

- 20 -

Soil removal performance on knitted cotton and polyviscose fabrics was measured in a tergotometer test. The soils used were

- 5 Soil A: soya bean oil (chosen as a typical greasy kitchen soil), coloured with a violet dye (0.08 wt%) to act as a visual indicator.

- 10 Soil B: paraffinic oil with particulate iron and carbon dispersed therein (no indicator dye was needed because the soil was itself sufficiently coloured by the presence of the particulate material)

Soil C: multi-use oil.

15

Test cloths (10 cm x 10 cm), each soiled with 0.5 ml of one of the soils listed above, were washed in tergotometers using the detergent compositions above under the following conditions:

20

Temperature	25°C
Liquor to cloth ratio	30:1
Product dosage	2.0 g/l
Soak time	10 min
Wash time (agitation)	15 min

The water used was of a hardness corresponding to a pK_{Ca}^{2+} of 6.4.

- 25 The reflectance ΔE , indicative of total colour change (of the violet dye) across the whole visible spectrum, of each

- 21 -

test cloth was measured before and after the wash. The results, expressed as the difference $\Delta\Delta E$ between reflectance values ΔE before and after the wash, are shown in the following tables.

5

Example 1, Comparative Examples A to C:

Soil A on knitted polyviscose

Example	wt% of total surfactant			$\Delta\Delta E$
	LAS	Nonionic	Cationic	
A	100	0	0	8.6
B	90	10	0	9.9
C	90	0	10	8.7
1	90	5	5	10.7

10

Examples 2 and 3, Comparative Examples D to F:

Soil C on knitted cotton

Example	wt% of total surfactant			$\Delta\Delta E$
	LAS	Nonionic	Cationic	
D	100	0	0	20.0
E	80	20	0	15.5
F	80	0	20	19.3
2	80	10	10	21.6
3	60	20	20	22.5

- 22 -

Examples 4 and 5, Comparative Examples G to I:Soil C on knitted polyviscose

Example	wt% of total surfactant			$\Delta\Delta E$
	LAS	Nonionic	Cationic	
G	100	0	0	16.0
H	80	20	0	2.2
I	80	0	20	14.0
4	80	10	10	18.8
5	60	20	20	17.8

5

Examples 6 to 8, Comparative Examples J to N:Soil B on knitted polyviscose

Example	wt% of total surfactant			$\Delta\Delta E$
	LAS	Nonionic	Cationic	
J	100	0	0	3.64
K	90	10	0	4.9
L	90	0	10	3.4
6	90	5	5	8.0
M	80	20	0	3.6
N	80	0	20	3.1
7	80	10	10	4.7
8	60	20	20	6.0

10

- 23 -

EXAMPLE 9, COMPARATIVE EXAMPLES P and Q:


performance appraisal of anionic/nonionic/cationic
surfactant mixtures containing ethoxylated cationic
surfactant (DBETAC)

5

High-suds detergent compositions according to the general
 formulation given in the previous Examples were prepared
 containing the surfactant systems shown below. The total
 amount of surfactant in each case was 24 wt% of the
 10 formulation.

DBETAC is the compound of the formula V as given previously:

15



20 wherein R₉ is a C₁₂-C₁₄ alkyl group.

Example	LAS	Nonionic AO30	Cationic DBETAC
P	100	0	0
Q	50	0	50
9	40	20	20

25 Performance was appraised, using the method of previous
 Examples, on Soil B and knitted cotton or polyviscose in
 water of three different hardnesses.

- 24 -

Knitted Cotton

	Water hardness		
	6° French	20° French	40° French
P	13.0	10.5	13.9
Q	8.2	5.1	5.8
9	13.9	14.4	12.2

5 Knitted polyviscose

	Water hardness		
	6° French	20° French	40° French
P	4.7	9.3	9.2
Q	5.0	4.0	1.8
9	11.3	11.6	9.7

- 10 These results show how the ternary system gives a robust performance across a wide range of water hardnesses.

- 25 -

EXAMPLES 10 to 12: PARTICULATE DETERGENT COMPOSITIONS
CONTAINING LUTENSOL AO30 AND PRAEPAGEN HY

Example 10

- 5 High suds formulation similar to that used in earlier Examples

LAS	21.60
Nonionic (Lutensol AO30)	1.20
Cationic (Praepagen HY)	1.20
Total surfactant	24.00
Sodium tripolyphosphate	19.00
Sodium carboxymethyl cellulose	0.33
Sodium neutral silicate	6.98
Sodium sulphate	13.70
Fluorescers	0.19
Acrylic/maleic copolymer	1.50
Sodium carbonate	15.00
Sodium perborate monohydrate	8.00
Tetracetyl ethylenediamine	2.40
Phosphonate sequestrant	0.40
Enzyme granules	0.91
Soil release polymer	0.80
Perfume	0.30
Miscellaneous salts, water etc	to 100

- 26 -

Example 11

Low suds formulation suitable for use in a closed drum
washing machine

LAS	5.80
Nonionic (Lutensol AO30)	2.00
Cationic (Praepagen HY)	2.00
Total surfactant	9.80
Soap	4.00
Sodium tripolyphosphate	25.00
Sodium carboxymethyl cellulose	0.50
Sodium neutral silicate	8.96
Sodium sulphate	22.84
Fluorescers	0.13
Sodium carbonate	6.31
Sodium perborate monohydrate	5.84
Tetracetyl ethylenediamine	2.10
Phosphonate sequestrant	0.50
Enzyme granules	0.97
Antifoam granules	2.00
Soil release polymer	0.50
Perfume	0.36
Miscellaneous salts, water etc	to 100

- 27 -

Example 12

Medium suds formulation suitable for use in a top-loading washing machine or for washing by hand

LAS	13.00
Nonionic (Lutensol AO30)	2.40
Cationic (Praepagen HY)	0.50
Total surfactant	15.90
Sodium tripolyphosphate	34.00
Sodium carboxymethyl cellulose	0.50
Sodium silicate	7.00
Sodium hydroxide	0.45
Sodium chloride	2.00
Fluorescers	0.15
Silicone fluid antifoam	0.05
Acrylic polymer	1.00
Sodium aluminosilicate	0.50
Sodium carbonate	3.58
Sodium perborate tetrahydrate	7.67
Tetracetyl ethylenediamine	2.21
Enzyme granules	1.64
Soil release polymer	0.35
Citric acid	1.00
Antifoam granules	3.00
Coloured speckles (sodium tripolyphosphate)	1.80
Perfume	0.33
Miscellaneous salts, water etc	to 100

- 28 -

EXAMPLES 13 and 14: DETERGENT COMPOSITIONS CONTAINING
LUTENSOL AO30 AND DBETAC

Example 13

- 5 Low suds formulation suitable for use in a closed drum washing machine.

LAS	3.90
Nonionic (Lutensol AO30)	2.00
Cationic (DBETAC)	2.00
Total surfactant	7.90
Soap	4.00
Sodium tripolyphosphate	25.00
Sodium carboxymethyl cellulose	0.50
Sodium neutral silicate	8.96
Sodium sulphate	22.84
Fluorescers	0.13
Acrylic/maleic copolymer	-
Sodium carbonate	6.31
Sodium perborate monohydrate	5.84
Tetracetyl ethylenediamine	2.10
Phosphonate sequestrant	0.50
Enzyme granules	0.97
Antifoam granules	2.00
Soil release polymer	0.50
Perfume	0.36
Miscellaneous salts, water etc	to 100

- 29 -

Example 14

Medium suds formulation suitable for use in a top-loading washing machine or for washing by hand.

LAS	5.40
Nonionic (Lutensol AO30)	3.20
Cationic (DBETAC)	6.40
Total surfactant	15.00
Sodium tripolyphosphate	34.00
Sodium carboxymethyl cellulose	0.50
Sodium silicate	7.00
Sodium hydroxide	0.45
Sodium chloride	2.00
Fluorescers	0.15
Silicone fluid antifoam	0.05
Acrylic polymer	1.00
Sodium aluminosilicate	0.50
Sodium carbonate	3.58
Sodium perborate tetrahydrate	7.67
Tetracetyl ethylenediamine	2.21
Enzyme granules	1.64
Soil release polymer	0.35
Citric acid	1.00
Antifoam granules	3.00
Coloured speckles (sodium tripolyphosphate)	1.80
Perfume	0.33
Miscellaneous salts, water etc	to 100

- 30 -

CLAIMS

1 A built laundry detergent composition comprising

5

(i) from 5 to 40 wt%, preferably from 7 to 30 wt%, of
surfactant,

(ii) from 10 to 80 wt% of detergency builder,

10

(iii) optionally other detergent ingredients to 100 wt%,


characterised in that the surfactant (i) consists
essentially of:

15

(i) (a) an anionic sulphonate or sulphate surfactant,

(i) (b) an ethoxylated alcohol nonionic surfactant of
the general formula I

20



characterised in that R is a hydrocarbyl chain having
from 8 to 16 carbon atoms, and the average degree of
ethoxylation n is from 20 to 50,

25

(i) (c) a quaternary ammonium cationic surfactant.

30

- 31 -

2 A detergent composition as claimed in claim 1,
characterised in that the surfactant (i) consists
essentially of:

5 (i) (a) from 20 to 98 wt% of the anionic sulphonate or
sulphate detergent;

(i) (b) from 1 to 60 wt% of the nonionic surfactant, and

10 (i) (c) from 1 to 60 wt% of the cationic surfactant.

3 A detergent composition as claimed in any preceding
claim, characterised in that it comprises, based on the
15 composition:

(i) (a) from 1 to 20 wt% of the anionic sulphonate or
sulphate detergent;

20 (i) (b) from 0.5 to 20 wt% of the nonionic surfactant, and


(i) (c) from 0.1 to 20 wt% of the cationic surfactant.

25 4 A detergent composition as claimed in any preceding
claim, characterised in that the weight ratios between the
surfactants are within the following ranges:

(i) (a) : (i) (b)	1.5:1 - 25:1
(i) (a) : (i) (c)	0.2:1 - 5:1
(i) (b) : (i) (c)	0.1:1 - 3:1

- 32 -

5 A detergent composition as claimed in any preceding
claim, characterised in that the cationic surfactant is a
5 compound of the formula III:



15 in which R_1 is a C_8 - C_{18} alkyl group, more preferably a C_8 - C_{10}
or C_{12} - C_{14} alkyl group, R_2 is a methyl group, and R_3 and R_4 ,
which may be the same or different, are methyl or
hydroxyethyl groups.

6 A detergent composition as claimed in claim 5,
20 characterised in that the surfactant (i) consists
essentially of:

(i) (a) from 50 to 98 wt%, preferably from 60 to 95 wt%,
of the anionic sulphonate or sulphate detergent;

25

(i) (b) from 1 to 30 wt%, preferably from 5 to 25 wt%, of
the nonionic surfactant, and

(i) (c) from 1 to 30 wt% , preferably from 5 to 25 wt%, of
30 the cationic surfactant.

- 33 -

7 A detergent composition as claimed in claim 5 or claim 6, characterised in that it comprises, based on the composition:

5 (i) (a) from 3 to 30 wt%, preferably from 5 to 25 wt%, of the anionic sulphonate or sulphate detergent;

(i) (b) from 0.5 to 10 wt%, preferably from 1 to 5 wt%, of the nonionic surfactant, and

10

(i) (c) from 0.1 to 10 wt%, preferably from 0.2 to 5 wt%, of the cationic surfactant.

15 8 A detergent composition as claimed in any one of claims 5 to 7, characterised in that the weight ratios between the surfactants are within the following ranges:


		preferably
(i) (a) : (i) (b)	2:1 - 25:1	3:1 - 20:1
(i) (a) : (i) (c)	2:1 - 50:1	3:1 - 30:1
(i) (b) : (i) (c)	0.1:1 - 3:1	0.5:1 - 2:1

20

- 34 -

9 A detergent composition as claimed in any one of claims
1 to 4, characterised in that the cationic surfactant (i) (c)
is an ethoxylated quaternary ammonium compound of the
formula IV:

5



10

wherein R_5 is a C_6 - C_{20} alkyl group,
 m is an integer from 1 to 20,
 R_6 and R_7 , which may be the same or different, each
15 represents a C_1 - C_4 alkyl group or a C_2 - C_4
hydroxyalkyl group,
 R_8 represents a C_1 - C_4 alkyl group, and
 Y^- represents a monovalent solubilising anion.

20


10 A detergent composition as claimed in claim 9,
characterised in that the cationic surfactant (i) (c) is a
compound of the formula IV in which

25

R_5 is a C_{10} - C_{16} alkyl group,
 m is from 1 to 4,
 R_6 , R_7 and R_8 are methyl groups, and
 Y^- represents Cl^- .

- 35 -

11 A detergent composition as claimed in claim 10, characterised in that the cationic surfactant (i) (c) is a compound of the formula V:



10

wherein R₉ is a C₁₂-C₁₄ alkyl group.

12 A detergent composition as claimed in any one of claims
 15 9 to 11, characterised in that the surfactant (i) consists essentially of:

(i) (a) from 20 to 80 wt%, preferably from 30 to 60 wt%,
 of the anionic sulphonate or sulphate detergent;

20

(i) (b) from 5 to 40 wt%, preferably from 10 to 30 wt%, of
 the nonionic surfactant, and

(i) (c) from 10 to 60 wt% , preferably from 20 to 50 wt%,
 25 of the cationic surfactant.

- 36 -

13 A detergent composition as claimed in any one of claims 9 to 12, characterised in that it comprises, based on the composition:

5 (i) (a) from 3 to 30 wt%, preferably from 5 to 25 wt%, of the anionic sulphonate or sulphate detergent;

(i) (b) from 0.5 to 10 wt%, preferably from 1 to 5 wt%, of the nonionic surfactant, and

10

(i) (c) from 0.1 to 10 wt%, preferably from 0.2 to 5 wt%, of the cationic surfactant.

15 14 A detergent composition as claimed in any one of claims 9 to 13, characterised in that the weight ratios between the surfactants are within the following ranges:

		preferably
(i) (a) : (i) (b)	1.5:1 - 10:1	1:1 - 5:1
(i) (a) : (i) (c)	0.2:1 - 5:1	0.5:1 - 3:1
(i) (b) : (i) (c)	0.2:1 - 5:1	0.5:1 - 3:1

20

15 A detergent composition as claimed in any preceding claim, characterised in that the ethoxylated nonionic surfactant (i) (b) has a hydrocarbon chain containing from 10
25 to 16 carbon atoms, preferably from 12 to 15 carbon atoms.

- 37 -

16 A detergent composition as claimed in any preceding claim, characterised in that the ethoxylated nonionic surfactant (i) (b) has an average degree of ethoxylation n of from 25 to 40.

5

17 A detergent composition as claimed in claim 15 and claim 16, characterised in that the ethoxylated nonionic surfactant (i) (b) has a hydrocarbon chain containing from 10 to 16 carbon atoms and an average degree of ethoxylation n of from 25 to 40.

18 A detergent composition as claimed in any preceding claim, characterised in that the anionic surfactant (i) (a) is linear alkylbenzene sulphonate.

19 A detergent composition as claimed in any preceding claim, characterised in that the composition is free from nonionic surfactants other than the nonionic surfactant (i) (b).

20 A detergent composition as claimed in any preceding claim, characterised in that it comprises from 10 to 40 wt% of a detergency builder (ii) selected from sodium tripolyphosphate, zeolite, sodium carbonate, sodium citrate, layered silicate, and combinations thereof.

30

- 38 -

21 A detergent composition as claimed in any preceding
claim, characterised in that it comprises one or more
optional ingredients (iii) selected from soap, peroxyacid
and persalt bleaches, bleach activators, sequestrants,
5 cellulose ethers and esters, other antiredeposition agents,
sodium sulphate, sodium silicate, sodium chloride, calcium
chloride, sodium bicarbonate, other inorganic salts,
fluorescers, photobleaches, polyvinyl pyrrolidone, other dye
transfer inhibiting polymers, foam controllers, foam
10 boosters, acrylic and acrylic/maleic polymers, proteases,
lipases, cellulases, amylases, other detergent enzymes,
citric acid, soil release polymers, fabric conditioning
compounds, coloured speckles, and perfume.

15

22 A detergent composition as claimed in any preceding
claim, characterised in that it is in powder form.

20 23 A process for laundering textile fabrics by machine or
hand, which includes the step of immersing the fabrics in a
wash liquor comprising water in which a laundry detergent
composition as claimed in any preceding claim is dissolved
or dispersed.


25

- 39 -

24 Use of a surfactant (i) consisting essentially of

(i) (a) an anionic sulphonate or sulphate surfactant,

5 (i) (b) an ethoxylated alcohol nonionic surfactant of
the general formula I



10 characterised in that R is a hydrocarbyl chain having
from 8 to 16 carbon atoms, and the average degree of
ethoxylation n is from 20 to 50,

(i) (c) a quaternary ammonium cationic surfactant,

15

in a laundry detergent composition in an amount of from 5 to
40 wt%, to improve the oily soil detergency of the
composition.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 01/14488

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D1/86 //C11D1:14,C11D1:22,C11D1:62,C11D1:72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 17769 A (PROCTER & GAMBLE) 30 April 1998 (1998-04-30) claims 1,11,12,15,18 page 1, line 3 page 10, line 7 - line 13 page 15, line 3 - line 7 example 4	1-5, 15-24
X	EP 0 000 225 A (PROCTER & GAMBLE) 10 January 1979 (1979-01-10) claim 1 page 14, line 19 -page 15, line 7 page 16, line 26 - line 29	1-4,9-24

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

25 April 2002

Date of mailing of the international search report

06/05/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Rasmusson, R

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/14488

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9817769	A	30-04-1998	GB 2318363 A	22-04-1998
			GB 2318362 A	22-04-1998
			GB 2323374 A	23-09-1998
			AT 213765 T	15-03-2002
			AU 4666897 A	15-05-1998
			AU 4805597 A	15-05-1998
			BR 9710446 A	17-08-1999
			BR 9712360 A	19-06-2001
			BR 9712363 A	31-08-1999
			BR 9712536 A	15-01-2002
			BR 9712541 A	24-10-2000
			BR 9712991 A	18-12-2001
			BR 9713260 A	19-06-2001
			BR 9713320 A	25-01-2000
			CN 1225669 A	11-08-1999
			CZ 9901352 A3	11-08-1999
			DE 69710749 D1	04-04-2002
			EP 0934397 A1	11-08-1999
			EP 0934388 A1	11-08-1999
			EP 0934389 A1	11-08-1999
			EP 0934377 A1	11-08-1999
			EP 0970169 A1	12-01-2000
			EP 0934379 A1	11-08-1999
			EP 0934378 A1	11-08-1999
			EP 0906386 A1	07-04-1999
			EP 0934391 A1	11-08-1999
			EP 0968269 A1	05-01-2000
			HU 0000115 A2	28-06-2000
			HU 0000117 A2	28-06-2000
			HU 0000118 A2	28-06-2000
			JP 2000504066 T	04-04-2000
			JP 2000504067 T	04-04-2000
			JP 2002509560 T	26-03-2002
			NO 991735 A	17-06-1999
			TR 9900851 T2	21-07-1999
			WO 9743367 A1	20-11-1997
			WO 9817751 A1	30-04-1998
			WO 9817752 A1	30-04-1998
			WO 9817753 A1	30-04-1998
			WO 9817777 A1	30-04-1998
			WO 9817754 A1	30-04-1998
			WO 9817755 A1	30-04-1998
			WO 9817759 A1	30-04-1998
			WO 9817766 A1	30-04-1998
			WO 9817767 A1	30-04-1998
			WO 9817768 A1	30-04-1998
			WO 9817769 A1	30-04-1998
			WO 9817758 A1	30-04-1998
			US 6103685 A	15-08-2000
			US 6087314 A	11-07-2000
EP 0000225	A	10-01-1979	AT 400244 B	27-11-1995
			AT 469178 A	15-12-1986
			AU 525063 B2	21-10-1982
			AU 3758778 A	03-01-1980
			BE 7 T1	07-12-1979
			BE 8 T1	07-12-1979
			BR 7804162 A	20-02-1979

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/14488

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0000225	A	BR 7804163 A	20-02-1979
		CA 1109759 A1	29-09-1981
		CA 1129751 A1	17-08-1982
		CH 643881 A5	29-06-1984
		DE 2857161 A1	06-03-1980
		DE 2857163 A1	06-03-1980
		EP 0000224 A1	10-01-1979
		EP 0000225 A1	10-01-1979
		FR 2423534 A1	16-11-1979
		FR 2423537 A1	16-11-1979
		GB 2040986 A ,B	03-09-1980
		GB 2040987 A ,B	03-09-1980
		GR 63770 A1	15-12-1979
		IT 1097064 B	26-08-1985
		JP 54039416 A	26-03-1979
		JP 54039417 A	26-03-1979
		MX 147560 A	14-12-1982
		MX 147454 A	03-12-1982
		NL 7815014 A	31-10-1979
		NL 7815014 T	31-10-1979
		NL 7815015 A	31-10-1979
		NL 7815015 T	31-10-1979
		PH 15326 A	24-11-1982
		SE 446460 B	15-09-1986
		SE 7907257 A	31-08-1979
		US 4321165 A	23-03-1982
		US 4333862 A	08-06-1982

